


Product no **AS11 1789****FtsH1-11 | ATP-dependent zinc metalloprotease FtsH1-11****Product information**

Immunogen	KLH-conjugated peptide derived from sequences of all known FtsH isoforms of <i>Arabidopsis thaliana</i> : FtsH1 Q39102 (At1g50250), FtsH2 Q80860 (At2g30950), FtsH3 Q84WU8 (At2g29080), FtsH4 Q80983 (At2g26140), FtsH5 Q9FH02 (At5g42270), FtsH6 Q1PDW5 (At5g15250), FtsH7 Q9SD67 (At3g47060), FtsH8 Q8W585 (At1g06430), FtsH9 Q9FIM2 (At5g58870), FtsH10 Q8VZ18 (At1g07510), FtsH11 Q9FGM0 (At5g53170) as well as 4 FtsH isoforms of <i>Synechocystis</i> sp. PCC6803NP_440330.1 , NP_442160.1 , NP_440797.1 , NP_440525.1
Host	Rabbit
Clonality	Polyclonal
Purity	Serum
Format	Lyophilized
Quantity	50 µl
Reconstitution	For reconstitution add 50 µl of sterile water
Storage	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

Application information


Recommended dilution	1 : 500-1 : 2000 (WB)
Expected apparent MW	<i>A. thaliana</i> : FtsH1:76.7, FtsH2: 74, FtsH5: 75.2, FtsH6: 74.5, FtsH7: 87.8, FtsH8: 73, FtsH9:87.8 kDa (chloroplastic); FtsH3: 89.3, FtsH4: 77.2, FtsH10: 89.5 kDa, FtsH11:88.7 kDa (mitochondrial)
Confirmed reactivity	<i>Acetabularia acetabulum</i> , <i>Arabidopsis thaliana</i> , <i>Bryopsis hypnoides</i> (ulvophytes), <i>Guillardia theta</i> , <i>Hordeum vulgare</i> , <i>Synechocystis</i> sp., <i>Thalassiosira pseudonana</i> , <i>Thalassiosira punctigera</i> , <i>Zea mays</i>
Predicted reactivity	Diatoms including <i>Phaeodactylum tricornutum</i> , dicots, <i>Chlamydomonas reinhardtii</i> , Monocots; <i>Nannochloropsis gaditana</i> , <i>Solanum lycopersicum</i>
	Species of your interest not listed? Contact us
Not reactive in	No confirmed exceptions from predicted reactivity are currently known
Additional information	For detection on a diatom samples, load of 5-10 µg/well is required as well as usage of enhanced chemiluminescence. The antibody also detects the chloroplast-encoded FtsH from <i>Thalassiosira pseudonana</i> (diatom) along with 3 related isoforms encode in the nuclear genome of <i>Thalassiosira pseudonana</i> . For detection in BlueNative, recommended dilution is 1: 500.
Selected references	Dogra et al. (2019). Oxidative post-translational modification of EXECUTER1 is required for singlet oxygen sensing in plastids. Nat Commun. 2019 Jun 27;10(1):2834. doi: 10.1038/s41467-019-10760-6. Li et al. (2016). A Hard Day's Night: Diatoms Continue Recycling Photosystem II in the Dark. Front. Mar. Sci., 08 November 2016 Tietz et al. (2015). Functional Implications of Photosystem II Crystal Formation in Photosynthetic Membranes. J Biol Chem. 2015 Apr 20. pii: jbc.M114.619841. Campbell et al. (2013). Photosystem II protein clearance and FtsH function in the diatom <i>Thalassiosira pseudonana</i> . Photosynth. Res. March 16.

Application example



8 µg of total protein from *Arabidopsis thaliana* wild type (1) and FtsH 1 and 3 deletion mutant (2) extracted with 0.2M Tris-HCl pH6.8, 2%SDS, 10% mercaptoethanol, 5M Urea were separated on 12 % **SDS-PAGE** and blotted 2h to **NC**. Blots were blocked with non fat milk for 1h at room temperature (RT) with agitation. Blot was incubated in the primary antibody at a dilution of 1: 2 000 over night at 4°C with agitation. The antibody solution was decanted and the blot was rinsed briefly twice, then washed once for 5 min and 2 times for 10 min in TBS-T at RT with agitation. Blot was incubated in secondary antibody anti-rabbit IgG horse radish peroxidase conjugated) diluted to 1:10 000 in TBS-T+5% non fat milk for 2h at RT with agitation. The blot was washed as above and developed for 5 min with ECL according to the manufacturers instructions. Exposure time was 3 minutes.

Courtesy of Dr. Leah Nave, The Hebrew University, Israel



The dilution series of *Hordeum vulgare* (4; 2; 1 µg Chl) and *Arabidopsis thaliana* (3,6; 1,8; 0,9 µg Chl) thylakoids were separated on 12% Criterion XT Bis-Tris SDS-PAGE (BioRad) gels and blotted for 25min 100V to PVDF membrane. Blot was blocked with 5% fat free skimmed milk in PBS-T for 1h at room temperature (RT) with agitation. Blot was incubated in the primary antibody at a dilution of 1: 500 overnight with agitation in 4°C. The antibody solution was decanted and the blot was washed 3 times for 5 min in PBS-T at RT with agitation. Blot was incubated in secondary antibody (swine anti-rabbit IgG horse radish peroxidase conjugated) diluted to 1: 5000 in 1% fat free skimmed milk in PBS-T for 1h at RT with agitation. The blot was washed 5 min in PBS-T and 1 min in PBS developed for 5 min with ECL according to the manufacturers instructions. Exposure time was 90 seconds.

The band below FtsH2 is suspected to be FtsH5 as according to Sinvány-Villalobo et al. (2004): The most abundant FtsH transcript is that of FtsH2 (Fig. 3B), demonstrating a level similar to ClpC1. FtsH1 and 5 are also relatively abundant, each accumulating to levels of about 50% to 60% of FtsH2 transcript. Other FtsH transcripts are much less abundant, with FtsH6 not accumulating at all under optimal growth conditions.

Courtesy of Dr. Marta Powikrowska, University of Copenhagen, Denmark