

Product no **AS16 3246****Goat anti-Bovine IgG (H&L), ALP conjugated****Product information**

Immunogen	Purified Bovine IgG, whole molecule
Host	Goat
Clonality	Polyclonal
Purity	Immunogen affinity purified goat IgG.
Format	Liquid
Quantity	1,0 mg
Storage	Store non-diluted antibody at 2-8°C. For storage at -20 °C, dilute with an equal volume of glycerol to prevent loss of enzymatic activity. Prepare working dilutions prior to use and then discard.

Additional information	<p>Antibody purity is > 95% based on SDS-PAGE. Antibody concentration is 1.5 mg/ml (E 1% at 280 nm = 13.0). Affinity purified using solid phase Bovine IgG (H&L). Antibody is supplied in 30 mM Triethanolamine, pH 7.2, 5 mM Magnesium Chloride, 0.1 mM Zinc Chloride, 1 % (w/v) BSA, Protease/IgG free. 0.05% (w/v) Sodium Azide is added as preservative.</p> <p>Based on IEP, this antibody reacts with: · heavy () chains on bovine IgG · light chains on all bovine immunoglobulins. Based on IEP, no reactivity is observed to: · non-immunoglobulin bovine serum immunoglobulins.</p>
-------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Application information

Recommended dilution	1 : 50 000-1 : 5 000 (ELISA), 1 : 500-1 : 2000 (IHC), 1 : 50 000-1 : 5 000 (WB)
Confirmed reactivity	Bovine IgG (H&L)