

Product no **AS16 3244****Goat anti-Bovine IgG (H&L), Unconjugated****Product information**

<b>Immunogen</b>	Purified Bovine IgG, whole molecule
<b>Host</b>	Goat
<b>Clonality</b>	Polyclonal
<b>Purity</b>	Immunogen affinity purified in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2 with 0.05% (w/v) sodium azide.
<b>Format</b>	Liquid
<b>Quantity</b>	2,0 mg
<b>Storage</b>	Store at 2-8°C.

<b>Additional information</b>	<p>Antibody purity is &gt; 95% based on SDS-PAGE          Affinity purified using solid phase Bovine IgG (H&amp;L)          Antibodi is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2.          0.05% (w/v) Sodium Azide is added as preservative.</p> <p>Based on IEP, this antibody reacts with: · heavy ( ) chains on bovine IgG · light chains on all bovine immunoglobulins.</p> <p>Based on IEP, no reactivity is observed to: · non-immunoglobulin bovine serum immunoglobulins.</p>
-------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

**Application information**

<b>Recommended dilution</b>	The optimal working dilution should be determined by the investigator
<b>Confirmed reactivity</b>	Bovine IgG (H&L)
<b>Selected references</b>	<a href="#">Patathil et al. (2010)</a> . A comprehensive toolkit of plant cell wall glycan-directed monoclonal antibodies. <i>Plant Physiol.</i> 2010 Jun;153(2):514-25.doi: 10.1104/pp.109.151985.