

Product no **AS10 1428**

## Goat anti-Mouse IgG (H&L), TRITC conjugated, min, cross-reactivity to bovine, horse, human, pig or rabbit serum

### Product information

**Immunogen** | Purified mouse IgG (H&L) [AAA51107](#)

**Host** | Goat

**Clonality** | Polyclonal

**Purity** | Immunogen affinity purified goat IgG.

**Format** | Lyophilized

**Quantity** | 1 mg

**Reconstitution** | For reconstitution add 1,1 ml of sterile water, Let it stand 30 minutes at room temperature to dissolve, Prepare fresh working dilutions daily

**Storage** | Store lyophilized material at 2-8°C. For long time storage after reconstitution, dilute the antibody solution with glycerol to a final concentration of 50% glycerol and store as liquid at -20°C, to prevent loss of enzymatic activity. For example, if you have reconstituted 1 mg of antibody in 1,1 ml of sterile water add 1,1 ml of glycerol. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard, Be sure to mix well but without foaming.

**Additional information** | Purity of this preparation is > 95% based on SDS-PAGE. Antibody concentration is 1.0 mg/ml (E 1% at 280 nm = 13.0). Antibody is supplied in 10 mM sodium phosphate, 0.15 M sodium chloride, pH 7.2.1 % (w/v) B, Protease/IgG free. Contains 0.05% (w/v) sodium azide as preservative of bacterial growth.

Based on immunoelectrophoresis, this antibody reacts with:heavy chains on mouse IgG, light chains on all mouse immunoglobulins. Based on immunoelectrophoresis, no reactivity is observed to: non-immunoglobulin mouse serum proteins, serum proteins from bovine, horse, human, pig, or rabbit.

### Application information

**Recommended dilution** | The optimal working dilution should be determined by the investigator, Suggested starting dilution(s): 1 : 20-1 : 2000 for most applications