

Product no [AS12 2337](#)**Goat anti-Rabbit IgG (H&L) -F(ab)'2 fragment, DyLight® 650 conjugated, min, cross-reactivity to bovine, human, or Mouse IgG or serum proteins****Product information****Immunogen** | Purified Rabbit IgG, whole molecule**Host** | Goat**Clonality** | Polyclonal**Purity** | Immunogen affinity purified goat IgG.**Format** | Lyophilized**Quantity** | 0,5 mg**Reconstitution** | For reconstitution add 0,55 ml of sterile water, Let it stand 30 minutes at room temperature to dissolve, Prepare fresh working dilutions daily**Storage** | Store lyophilized material at 2-8°C. Product is stable for 4 weeks at 2-8°C after rehydration. For long time storage after reconstitution, dilute the antibody solution with glycerol to a final concentration of 50% glycerol and store as liquid at -20°C, to prevent loss of enzymatic activity. For example, if you have reconstituted 0,5 mg of antibody in 0,55 ml of sterile water add 0,55 ml of glycerol. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard. Be sure to mix well but without foaming.**Additional information** | Conjugate is present in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free. 0.05 % (w/v) sodium azide is added as preservative.

Based on immunoelectrophoresis, this antibody reacts with: heavy () chains on rabbit IgG light chains on all rabbit immunoglobulins

No reactivity is observed to: non-immunoglobulin rabbit serum proteins IgG from bovine, human or mouse serum proteins from bovine, human or mouse

Application information**Recommended dilution** | 1 : 20-1 : 2000 for most applications**Not reactive in** |**Additional information** |