

Product no **AS10 1018****Goat anti-Rabbit IgG (H&L), F(ab)'2 fragment, TRITC conjugated**

## Product information

**Host** | Goat**Clonality** | Polyclonal**Purity** | Immunogen affinity purified IgG, F(ab)'2 fragment.**Format** | Liquid**Quantity** | 0,5 mg**Storage** | Store at 2-8°C. Shelf life of this product is one year from date of receipt.**Additional information** | This antibody reacts with the heavy chains on rabbit IgG and with the light chains on all rabbit immunoglobulins based on immunoelectrophoresis.

Minimum cross-reactivity is observed to non-immunoglobulin rabbit serum proteins based on immunoelectrophoresis.

Antibody is supplied in 10 mM sodium phosphate, 150 mM sodium chloride, pH 7.2, 1% (w/v) BSA, protease/IgG free and 0.05 % (w/v) sodium azide as preservative.

## Application information

**Recommended dilution** | The optimal working dilution should be determined by the investigator, (ICC), immunohistochemistry (IHC) (Flow cyt)**Selected references** | [Feng et al. \(2020\)](#). Intracellular expression of arginine deiminase activates the mitochondrial apoptosis pathway by inhibiting cytosolic ferritin and inducing chromatin autophagy. BMC Cancer. 2020 Jul 16;20(1):665.doi: 10.1186/s12885-020-07133-4.