

Product no **AS09 630****Rabbit anti-Goat IgG (H&L), ALP conjugated****Product information**

| | |
|-------------------------------|--|
| Immunogen | Purified Goat IgG, whole molecule |
| Host | Rabbit |
| Clonality | Polyclonal |
| Purity | Immunogen affinity purified rabbit IgG. |
| Format | Liquid |
| Quantity | 1 mg |
| Storage | Non-diluted antibody is stable for 4 years at 2-8°C. For storage at -20°C dilute antibody solution with an equal volume of glycerol to obtain final glycerol concentration of 50 % to prevent loss of enzymatic activity. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard, Be sure to mix well but without foaming. |
| Additional information | ALP conjugate is supplied in 30 mM Triethanolamine, pH 7,2, 5 mM Magnesium Chloride, 0,1 mM Zinc Chloride, 1 % (w/v) BSA, Protease/IgG free, 0,05 % (w/v) of sodium azide is added as preservative |

Application information

| | |
|-------------------------------|--|
| Recommended dilution | 1 : 500-1 :2000 (ELISA), 1 : 50-1 : 5000 (ICC), 1 : 20 -1 : 2000 (IHC), 1 : 500-1 :2000 (WB) |
| Confirmed reactivity | Goat IgG heavy and light chains (H&L) |
| Predicted reactivity | Goat IgG Heavy and Light chains (H&L) |
| Not reactive in | No confirmed exceptions from predicted reactivity are currently known |
| Additional information | No reactivity is observed to non-immunoglobulin goat serum proteins based in immunoelectrophoresis. BSA and milk have to be replaced by other blocking reagents, like donkey serum or commercial formulations which are free from bovine IgG. |