

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no AS22 4825

RbcS | Ribulose-1,5-bisphosphate carboxylase/oxygenase small subunit (Algal)

Product information

Immunogen KLH-conjugated peptide derived from Chlamydomonas reinhardti RbcS, UniProt: P00873, P08475, Q31NB2

Host Rabbit

Clonality Polyclonal

Purity Antigen affinity purified serum, in PBS pH 7.4

Format Lyophilized

Quantity 50 μg

Reconstitution For reconstitution, add 50 μl, of sterile or deionized water.

Storage Store lyophilized/reconstituted at -20 °C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please, remember to spin tubes briefly prior to opening them to avoid any losses that might occur from lyophilized

material adhering to the cap or sides of the tubes.

Application information

Recommended dilution 1:1000 (WB)

Expected | apparent

20 | 15 kDa

Confirmed reactivity Chlamydomonas reinhardtii, Synechococcus sp. PCC7002

Predicted reactivity | Species of your interest not listed? Contact us

Not reactive in No confirmed exceptions from predicted reactivity are currently known

Selected references To be added when available, antibody available in October 2023.



 $5~\mu g$ of total protein from *C.reinhardtii*,was denatured with Invitrogen LDS sample buffer (4X) at $70~^{\circ}$ C/5 min. Samples were separated on Invitrogen NuPage Bis-Tris 4-12% SDS-PAGE 60 min and blotted for 1 h to Invitrogen PVDF (pore size of 0.45 μm), using: wet transfer. Blot was blocked with 5% milk in TBS-T for: 1h/RT with agitation. Blot was incubated in the primary antibody at a dilution of 1: 1000 for 1h/RT with TBS-T Blocking. The antibody solution was decanted, and the blot was rinsed briefly twice, then washed once for 10 min and 2 times for 5 min in TBS-T at RT with agitation. Blot was incubated in matching secondary antibody (anti-rabbit IgG ALP conjugated, <u>AS09 607</u> lot 2302) diluted to 1: 5 000 in TBS-T Blocking for 0,5h/RT with agitation. The blot was washed as above and developed with <u>AS19 BCIP-NBT</u> lot 03068231 for 30 seconds. Image was captured after 1h.